



References

- [1] A. Baklanov, A. Aloyan, A. Mahura, V. Arutyunyan, and P. Luzan. Evaluation of source receptor relationship for atmospheric pollutants using approaches of trajectory modelling, cluster, probability fields analyses and adjoint equations. *Atmos. Poll. Res.*, 2(4):400–408, 2011.
- [2] T. Balk, J. Kukkonen, K. Karatzas, T. Bassoukos, and V. Epitropou. A european open access chemical weather forecasting portal. *Atmos. Environ.*, 45(38):6917 – 6922, 2011.
- [3] O. Hänninen, G. Hoek, E. Mallone, S. annd Chellini, C. Katsouyanni K. and, Gariazzo, G. Cattani, A. Marconi, P. Molnar, and T. Bellander. Seasonal patterns of outdoor pm infiltration into indoor environments: review and meta-analysis of available studies from different climatological zones in europe. *Air Quality, Atmos. and Health*, 4(3 – 4):221–233, 2011.
- [4] J. P. Jalkanen, L. Johansson, J. Kukkonen, A. Brink, J. Kalli, and T. Stipa. Extension of an assessment model of ship traffic exhaust emissions for particulate matter and carbon monoxide. *Atmos. Chem. Phys.*, 12:2641–2659, 2012.
- [5] N. Janssen, G. Hoek, M. Simic-Lawson, P. Fischer, L. Bree, H. Ten-Brink, M. Keuken, R. Atkinson, H. Anderson, B. Brunekreef, and F. Cassee. Black carbon as an additional indicator of the adverse health effects of airborne particles compared with pm10 and pm2.5. *Environ. Health Per.*, 19(12):1691–1699, 2011.
- [6] M. Kauhaniemi, J. Kukkonen, J. Härkönen, J. Nikmo, L. Kangas, G. Omstedt, M. Ketzel, A. Kousa, M. Haakana, and A. Karppinen. Evaluation of a road dust suspension model for predicting the concentrations of pm(10) in a street canyon. *Atmos. Environ.*, 45(22):3646 – 3654, 2011.
- [7] M. Keuken, H. D. van der Gon, and K. van der Valk. Non-exhaust emissions of pm and the efficiency of emission reduction by road sweeping and washing in the netherlands. *Sci. Total Environ.*, 408(20):4591 – 4599, 2010.
- [8] J. Kukkonen, T. Olsson, D. M. Schultz, A. Baklanov, T. Klein, A. I. Miranda, A. Monteiro, M. Hirtl, V. Tarvainen, M. Boy, V. H. Peuch, A. Poupkou, I. Kioutsioukis, S. Finardi, M. Sofiev, R. Sokhi, K. E. J. Lehtinen, K. Karatzas, R. San José, M. Astitha, G. Kallos, M. Schaap, E. Reimer, H. Jakobs, and K. Eben. A review of operational, regional-scale, chemical weather forecasting models in europe. *Atmos. Chem. Phys.*, 12(1):1 – 87, 2012.
- [9] J. Langner, M. Engardt, A. Baklanov, J. H. Christensen, M. Gauss, C. Geels, G. B. Hedegaard, R. Nuterman, D. Simpson, J. Soares, M. Sofiev, P. Wind, and Zakey A. A multi-model study of impacts of climate change on surface ozone in europe. *Atmos. Chem. Phys. Discuss.*, 12:4901 – 4939, 2012.

- [10] V. Penenko, A. Baklanov, E. Tsvetora, and A. Mahura. Direct and inverse problems in a variational concept of environmental modelling. *Pure Appl. Geophys.*, 169:447–465, 2012.
- [11] A. Vlachogianni, P. Kassomenos, Ari Karppinen, S. Karakitsios, and Jaakko Kukkonen. Evaluation of a multiple regression model for the forecasting of the concentrations of no(x) and pm(10) in athens and helsinki. *Sci. Total Environ.*, 409(8):1559 – 1571, 2011.
- [12] D. Voukantsis, K. Karatzas, J. Kukkonen, T. Räsänen, A. Karppinen, and M. Kolehmainen. Intercomparison of air quality data using principal component analysis, and forecasting of pm(10) and pm(2.5) concentrations using artificial neural networks, in thessaloniki and helsinki. *Sci. Total Environ.*, 409(7):1266 – 1276, 2011.